



## Origins of the performance gaps in innovative cropping systems under experimental assessment.

Caroline Colnenne-David, Gilles Grandeau, Véronique Tanneau, Marie-Helene Jeuffroy, Thierry Doré

### ► To cite this version:

Caroline Colnenne-David, Gilles Grandeau, Véronique Tanneau, Marie-Helene Jeuffroy, Thierry Doré. Origins of the performance gaps in innovative cropping systems under experimental assessment.. The 5th International Symposium for Farming Systems Design “Multi-functional farming systems in a changing world”, Sep 2015, Montpellier, France. hal-01357467

**HAL Id: hal-01357467**

**<https://hal-agroparistech.archives-ouvertes.fr/hal-01357467>**

Submitted on 30 Aug 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Context and Objectives

To meet new agricultural issues and make agriculture more sustainable, innovative cropping systems (ICSs) targeting a multiplicity of purposes need to be designed. Four ICSs were designed by prototyping (Reau and Doré, 2008<sup>1</sup>) and assessed in a long-term field system experiment.

**Our objective was to analyze the results after the first complete rotation, particularly the bad performances compared to the targeted aims.**

## Materials and Methods

### Multiple goals of the four designed ICSs (Colnenne-David and Doré, 2014<sup>2</sup>)

#### The PHEP ICS goals:

##### 1. To satisfy multiple environmental criteria:

\*low pesticide use → high crop diversity, highly resistant varieties

\*low direct energy consumption → only 1 ploughing within the rotation

\*low indirect energy consumption → legumes in the rotation

\*low nitrogen leaching → catch crop (CC) before spring crops and no N fertilization during autumn and winter

\*stabilization and/or to enrich soil organic matter → burying residues of all crops

##### 2. To reach yield targets

matching the Ile-de-France yields

**Crop sequence:** winter faba bean, winter wheat, winter oilseed rape, winter wheat, mustard as CC and spring barley

#### The L-GHG ICS goals:

##### 1. 50% GHG emissions compared to the PHEP ICS

(i) increase soil C sequestration → many cereals, continuous soil cover, high yield targets, no ploughing

(ii) decrease N<sub>2</sub>O emissions → high number of legume crops in the rotation, improvement of N fertilization management, crops with taproots in order to reduce soil compaction

##### 2. To satisfy multiple environmental criteria: idem PHEP ICS

##### 3. To reach yield targets

matching the Ile-de-France yields  
**Crop sequence:** catch crop (CC), maize, triticale, CC, spring faba bean, winter oilseed rape, winter wheat, CC, winter barley

#### The L-EN ICS goals:

##### 1. 50% fossil energy consumption compared to the PHEP ICS

(i) Low direct energy consumption → no ploughing and using direct sowing machine

(ii) Low indirect energy consumption → high number of legume species in the rotation, species with high N efficiency use, decrease N fertilization by reducing yield objectives

##### 2. To satisfy multiple environmental criteria: idem PHEP ICS

##### 3. To reach yield targets: 20% lower than the Ile-de-France yields

**Crop sequence:** winter faba bean, winter wheat, winter flax, winter wheat–trifolium mixture, Trifolium as CC, spring oat

#### The No-Pest ICS goals:

##### 1. No pesticide is allowed

→ long rotation including a wide diversity of species (e.g. hemp), alternate sowing dates, different dates and densities of sowing, highly resistant varieties or mixtures, ploughing and mechanical weeding


##### 2. To satisfy multiple environmental criteria: idem PHEP ICS

##### 3. To reach yield targets: higher than organic systems in the Ile-de-France

**Crop sequence:** triticale, CC, maize, winter wheat, CC, spring faba bean, winter wheat, CC, hemp

### Main characteristics of the field trial

- ✓ located in Grignon (78, France)
- ✓ 6.2 ha (surface plot: ≈ 4000 m<sup>2</sup>)
- ✓ 3 blocks
- ✓ deep loamy soil
- ✓ Beginning of field assessment: 2008



## Results: Classification of the major disparities

Classification	Examples collected in the ICSs
Some agronomical strategies were no suitable to reach the goals	In the L-GHG ICS: No ploughing → No increase of C sequestration as expected C sequestration evolution = -149kgCO <sub>2</sub> ha <sup>-1</sup> year <sup>-1</sup> (+87kgCO <sub>2</sub> ha <sup>-1</sup> year <sup>-1</sup> expected)
Some practices were not adapted to satisfy a multiplicity of objectives	In the No-Pest ICS: No possible to satisfy both the pesticide constraint and the S.O.M. criteria. Restitution of small organic matter amounts + regular ploughings → Few weeds but adverse effect on C sequestration (C sequestration evolution = -560kgCO <sub>2</sub> ha <sup>-1</sup> year <sup>-1</sup> )
Some practices were not appropriate in the context of the field-trial conditions	In the L-GHG ICS: Very dry conditions in summer 3 years / 6 → Low amount of aerial biomass of cover crops
An unpredicted evolution of the agrosystem occurred	In both the L-EN and the L-GHG ICSs: High weed development → to mow oilseed rape plots in 2014

## Discussions – Conclusion

- After the first complete rotation the major sources of disparities were classified
- Nevertheless, a more complete agronomic diagnosis is necessary to identify and to rank all the causes of bad performances
- This knowledge allowed us to improve the innovative cropping systems through a new design step
- This experiment contributes to the learning design processes and cropping system management

### REFERENCES

<sup>1</sup> Reau R. and Doré T. (2008). "Systèmes de culture innovants et durables : quelles méthodes pour les mettre au point et les évaluer ?" *Educagri Editions, Dijon, France*, pp 175

<sup>2</sup> Colnenne-David C. and Doré T., 2014. Designing innovative productive cropping systems with quantified and ambitious environmental goals. "Renewable Agriculture and Food Systems".

doi:10.1017/S1742170514000313.